


Naive Set Theory: Basic Concepts

September 2, 2008



notation: If a is a member of set Y , $a \in Y$. If all the elements of set X are included in set Y , $X \subseteq Y$.

extension axiom: $(X = Y) \leftrightarrow (\forall a)(a \in X \leftrightarrow a \in Y) \leftrightarrow (X \subseteq Y \wedge Y \subseteq X)$

some simple sets:

$$\emptyset = \{a \mid a \neq a\}$$

$$X \cup Y = \{a \mid a \in X \vee a \in Y\}$$

$$X \cap Y = \{a \mid a \in X \wedge a \in Y\}$$

$$X - Y = \{a \mid a \in X \wedge a \notin Y\}$$

more complicated sets:

$$\langle a, b \rangle = \{\{a\}, \{a, b\}\}$$

$$X_1 \times \dots \times X_n = \{\langle x_1, \dots, x_n \rangle \mid x_1 \in X_1 \wedge \dots \wedge x_n \in X_n\}$$

binary relation R on X is a subset of $X \times X$

function f from X to Y is a binary relation between X and Y s.t. $(\forall x \in X)(\exists \text{ unique } y \in Y)(x f y)$

equivalence relations:

(i) R is reflexive if $(\forall a \in X)(Raa)$

(ii) R is symmetric if $(\forall a, b \in X)(Rab \rightarrow Rba)$

(iii) R is transitive if $(\forall a, b, c \in X)((Rab \wedge Rbc) \rightarrow Rac)$

- an R that satisfies (i), (ii), and (iii) is an *equivalence relation*
- if $a \in X$, its *equivalence class* is defined as $[a] = \{b \mid b \in X \wedge Rab\}$